

# Facilitators of and Barriers to Physical Activity Promotion among Physical Therapists who Treat Clients with Neurological Conditions: A Qualitative Study Informed by the Theoretical Domains Framework

Winston Kennedy<sup>1</sup>, Carol Curtin<sup>2</sup>, Jafrā D. Thomas<sup>2</sup>, and April Bowling<sup>4</sup>

<sup>1</sup>Northeastern University;

<sup>2</sup>UMass Chan Medical School;

<sup>3</sup>California Polytechnic State University;

<sup>4</sup>Merrimack College

Healthcare provider communications about physical activity (PA) lead to meaningful and sustained increases in PA, but physical therapists (PTs) do not consistently promote PA for clients with neurological conditions (NC). The purpose of this qualitative study was to understand factors that could support implementation strategies related to PTs promoting PA for clients with NC. The Theoretical Domains Framework (TDF) was used to develop a semi-structured interview guide. One-on-one online interviews were conducted in 2022 with 10 physical therapists licensed to practice in the United States whose caseloads include clients with NC. Responses were first sorted into a priori factors derived from previous research (TDF codebook), then reappraised to ensure other factors were not overlooked (inductive analysis). Responses about facilitators corresponded to three factors: Personal (e.g., sense of responsibility, personal connections to community opportunities); Professional (e.g., a specific PT course on PA promotion); and Client-related (e.g., highly motivated). Barriers corresponded to two factors: Professional (e.g., limited continuing education, lack of guidance/communication from PT-associations), and Client-related (e.g., lack of low-cost/free services to offer, provider safety concerns). There are opportunities for PTs to increase PA promotion when treating clients with NC, as outlined in PT education. PTs should also utilize and advocate for community-based partnerships and expand education/resources from PT associations beyond traditional communication channels.

*Keywords:* health promotion, implementation, disability health

Physical therapists (PTs) are health professionals who may be able to include physical activity (PA) promotion for people with neurological conditions (NC); however, it is well documented that PTs do not consistently engage in PA promotion behaviors as part of their clinical practice (Kennedy et al., 2023; Rethorn et al., 2021).

Including PA promotion in clinical practice is not a novel recommendation but instead is an evidence-based strategy. One scoping review found that PA promotion by clinical providers had a small-to-moderate positive effect on increasing clients' PA levels (Sanchez et al., 2015). PA promotion in clinical settings has also been shown to be effective for children and adults with chronic disabilities, including those with NC (Ross et al., 2021; van der Ploeg et al., 2007).

Many people with disabilities have greater difficulty participating in various types of health-promoting behaviors like PA engagement, social activities, and accessing preventative care, which contribute to health disparities that promote people with disabilities being at increased risk for secondary health conditions (Rimmer et

---

The authors have no conflict of interest to disclose.

This work was funded under the Department of Education no. H325D160023.

Correspondence concerning this article should be addressed to Winston Kennedy, Bouve College of Health Sciences, Northeastern University, 360 Huntington Ave, 301R Robinson Hall, Boston, MA 02115. Email: wi.kennedy@northeastern.edu

al., 2014). Specifically, youth and adults with disabilities have higher rates of secondary health conditions like pain, fatigue, depression, and obesity (Havercamp & Scott, 2015; Kinne et al., 2004; Rimmer et al., 2011), and report lower rates of participation in health promoting activities that are intended to change behaviors like physical activity engagement (Carroll et al., 2014; Rimmer, 2012; Rimmer et al., 2014). This is important for people with NC because they may either identify as being disabled or may be identified as disabled by a healthcare professional. Barriers to participating in health promotion initiatives include lack of staff or professionals who understand how to support people with disabilities in their programs, and negative perceptions and attitude among health professionals that support the idea that people with disabilities need 'specialized' services and support that are independent of and separated from the general public (Anderson et al., 2017; Ginis et al., 2016). Given that PA is a major health indicator for people with NC for functional recovery and is a mechanism of preventive care for secondary conditions, it is also vital to provide PA promotion for people with NC.

The limited implementation of evidence-based practice behaviors, such as PA promotion, in healthcare is common. To encourage better adoption of evidence-based practices, it is important to identify key factors associated with PA promotion among PTs (French et al., 2012; Grol & Grimshaw, 2003). The Theoretical Domains Framework (TDF), developed in implementation science, aims to facilitate behavior change among health professionals (Cane et al., 2012). The TDF has been used to assess how often PTs promote PA for clients with and without NC. Kennedy et al. (2023) used a TDF-informed survey to explore PA promotion practices among PTs treating clients with NC, finding that although most PTs viewed PA promotion as part of their scope of practice, their actual promotion behaviors were inconsistent. Similarly, Rethorn et al. (2021) surveyed outpatient PTs and identified knowledge, skills, environmental context, and social influences as key TDF domains linked to attitudes and practices related to PA promotion. However, these quantitative studies did not thoroughly explore the detailed barriers and facilitators PTs face, especially when serving clients with NC. More recently, the TDF was used to identify strategies to support the implementation of guidelines for the physical therapy management of people with spinal cord injuries using qualitative methods (Tranter et al., 2025). The barriers to implementing the guidelines identified by PTs were a lack of knowledge and skills, a lack of resources, and challenges associated with working within large organizations. Facilitators of the uptake of care guidelines included education, skills training, awareness, and two-way communication be-

tween patient and therapist. These findings are on par with barrier and facilitator research on PA promotion in clinical care (Tranter et al., 2025). However, the context is different because of the management of acute health risks for individuals who have an acute spinal cord injury, compared to those with chronic NC who are engaging in general PA. The aforementioned study is an example of how qualitative inquiry using the TDF can provide a rich, contextual understanding of implementation challenges and opportunities that quantitative surveys cannot capture.

Much of the PA promotion literature involving PTs has focused on implementation with populations without disabilities. Barriers to PA promotion in clinical practice for the general population include time, provider reimbursement, training, provider self-efficacy, healthcare system support, and knowledge of community resources (Lobelo et al., 2018). When considering PT specifically, additional barriers include: perceived lack of interest or awareness on the part of clients, the public, and other healthcare professionals that PT can provide PA promotion services (Kearns et al., 2014; Taukobong et al., 2014; Verhagen & Engbers, 2008); limited capacity to implement evidenced-based practice (Dzewaltowski et al., 2004; Eccles et al., 2005; Grimshaw et al., 2012); lack of participation of PT in health promotion research, policy, and action initiatives; and payment systems that do not emphasize wellness through PA promotion (Dean, 2009).

While the existing literature enumerates potential barriers to PA promotion experienced by PTs, qualitative research exploring PT perception of barriers and facilitators is lacking. This is especially true in the context of services provided to clients with NC, given that most of the research has focused on clients without a chronic disability. The purpose of the present study was to gain insight into PTs' perceived barriers and facilitators to promoting PA when serving clients with NC. The findings of this study should provide a focus for future implementation strategies to increase PTs' likelihood of promoting PA when serving clients with NC.

## Methods

This study was approved by the Institutional Review Board at Oregon State University in 2022 (the primary investigator's prior institution). The first author conducted all interviews.

## Participants

Purposive sampling was used to recruit licensed PTs who worked with clients with NC in the US. The sampling occurred from the pool of PTs who participated in another

study ( $N = 76$ ) (Kennedy et al., 2023). As per best practices for theory-informed qualitative research, we attempted to recruit participants until theme saturation was achieved (Francis et al., 2010). Data saturation was considered achieved when no new information emerged from three consecutive interviews, resulting in 10 interviews (after the 7th interview). Demographic data are shown in Table 1.

## Measures

A semi-structured interview guide was developed based on a review of the TDF literature (Cane et al., 2012). To ensure quality and understandability, questions in the interview guide were reviewed by health promotion experts and PTs who were not part of this study. The expert review process involved interviews with three experts to assess the clarity, comprehension, and relevance of the questions to the research aims (Kallio et al., 2016). Updates to the interview guide were made based on the expert's feedback. The interview guide is shown in Table 4.

## Data Analysis

Interviews were conducted and recorded via Zoom™ (Archibald et al., 2019). Transcripts were downloaded, transcribed, cleaned, and read. Analysis of the transcriptions used a pragmatic approach that combined applicable evidence-based qualitative strategies (Ramanadhan et al., 2021). Thematic analysis was then used as the analytic strategy to represent the data through qualitative description (Sandelowski, 2000). Transcriptions were initially coded deductively using a TDF-based codebook (Atkins et al., 2017). Data were then re-analyzed inductively to ensure that important factors were not lost during the deductive data analysis (Fereday & Muir-Cochrane, 2006). The data were divided into facilitator and barrier categories to both align with and extend prior research on health professionals promoting health for their clients. All transcripts were analyzed by the first author (primary investigator) and then independently reviewed by the fourth author, ensuring a rigorous and trustworthy analysis. A third reviewer was available to resolve any discrepancies if necessary.

## Results

Identified trends were categorized into three groups of facilitators that facilitate PTs' implementation of PA promotion: personal, professional, and client-related. Two categories of barriers were identified: professional and client-related. Personal facilitators for PTs were related to participants' sense of self and to how that supported the promotion of PA. Professional facilitators for

PTs include factors that support PA promotion through PT training and professional practice. Client-related facilitators were client characteristics that PTs believed facilitated their ability to promote PA as part of their service to a client. See Table 2 for identified trends and examples. Professional barriers for PTs stemmed from factors in PT training and practice that made it harder to engage in PA promotion. Client-related barriers for PTs were client-specific factors (e.g., environmental resources, perceived risk of injury). See Table 3 for identified trends and examples.

## Discussion

The present study sought to understand how PTs perceived and experienced barriers to and facilitators of PA promotion when treating clients with NC based on the TDF. Facilitators and barriers for PTs to increase uptake in PA promotion in clinical practice that were identified may be able to inform implementation practices that can increase the uptake of including PA promotion by PTs who treat clients with NC.

Several personal facilitator factors were identified that may be understood through the lens of public service motivation (PSM). PSM refers to an individual's motivation to serve the public interest, encompassing beliefs, values, and attitudes that go beyond self-interest to concern the interests of a larger community (Perry & Wise, 1990). Importantly, research has shown that individuals with high levels of PSM are more likely to engage in behaviors that serve the public good, including health promotion activities in healthcare settings (Roh et al., 2016). This suggests that PSM may function as an important antecedent to PA promotion behaviors among PTs, particularly when serving populations with heightened health needs such as clients with NC. While PSM was originally conceptualized as specific to public-sector employment, research suggests that PSM manifests across sectors when professionals perform similar public-serving tasks (Andersen et al., 2011).

The personal facilitators identified in the current study appear to be manifestations of PSM dimensions that may increase the likelihood of PTs engaging in PA promotion for clients with NC. Participants' sense of responsibility to learn about PA promotion and their confidence in promoting PA align with the commitment-to-public-interest dimension of PSM, suggesting that PTs who feel a duty to serve the broader health needs of people with NC may be more inclined to incorporate PA promotion into their practice. Similarly, participants' reports of personal enjoyment and satisfaction from promoting PA reflect the compassion dimension of PSM, indicating that the intrinsic reward from helping clients with NC improve

**Table 1:** *Participant Demographics*

Characteristics	N (10)
<i>Gender</i>	
Male	2
Female	8
Average age, years (range)	44 (30–72)
Average experience, years (range)	18 (3–49)
<i>Race/Ethnicity</i>	
White (non-Hispanic/Latinx)	7
Asian	2
Other	1
<i>Highest Level of Education</i>	
Academic doctorate (DSc, EdD, PhD, etc.)	1
Bachelor's	2
Clinical doctorate (DPT, tDPT, etc.)	6
Master's	1
<i>Practice Setting</i>	
Home health	1
Commercial PT (5 or more locations) OP	1
Private Practice (PT-owned) OP	4
Neurological OP	2
Hospital-based OP	2
<i>Patient Age Range</i>	
<18	4
18–64	5
≥65	1

their health through PA may motivate continued engagement in these promotion behaviors. This interpretation is supported by a recent systematic review, which found that health professionals with strong PSM are more likely to choose and remain in positions that allow frequent contact and interaction with patients (Fernandes et al., 2022). The review further identified that altruistic PSM-related factors (desire to help patients, recognition of one's contribution to public health) sustain engagement in public service activities. This suggests that PTs who demonstrate high PSM, particularly in the compassion and public-interest dimensions, may be more likely to overcome barriers to PA promotion and to persist in PA promotion despite systemic challenges. Understanding that these personal facilitators are linked to PSM provides a theoretical foundation for implementation strategies related to strengthening PTs' sense of public service mission and their connection to the health outcomes of people with NC.

Among the multiple professional facilitator factors identified, having community connections for NC-specific resources indicates the impact community engagement can have on health professional practice. Participants speaking saliently of this professional facilitator was surprising, given its absence from PT education. Further, prior research suggests PTs are resistant to referring clients with NC to community-based PA programs (Williams et al., 2018). When drilling down into why PTs may resist referring clients to community-based programs, Williams

et al. (2018) found that PTs were concerned programs would promote unrealistic expectations or would be run unsafely. An implementation strategy suggested by our findings for increasing referrals to community-based PA promotion programs is for PTs to partner with community programs. Most participants in our study were connected to a community PA opportunity for individuals with chronic disabilities. This community PA involvement was a salient professional facilitator, meaning they may have seen it as an extension of their practice to reduce barriers to PA and support clients in engaging in safe forms of exercise and active recreation.

Within the professional facilitator category, participants reported that their PT education included relevant content on people with disabilities and that they had participated in continuing education relevant to PA promotion for clients with NC. This suggests that these PTs may have a better understanding of how to promote PA for their clients with NC; however, this warrants further investigation. Although this may not be true for all PT training curricula, there is a shift to include more content on people with disabilities (Garrand et al., 2018; Robey et al., 2013; Sharp & Herrman, 2021; Whalen Smith et al., 2024). Further, participants reported that their job helped them promote PA, that they worked at a hospital with community programs specifically focused on exercise, and that the organization in which a PT works has the potential to assist PTs in promoting PA. This is not a surprising finding because supportive environments

are conducive to health professionals' satisfaction, which may, in turn, support their work performance (Sumathi et al., 2015). It can be assumed that organizational support reduces the burden on the sole PT to promote PA, which may make it more likely that the PT will engage in the behavior.

Several professional barriers were identified. Acknowledging these barriers provides an opportunity to implement strategies to reduce them. Pivotal professional barrier factors included a lack of knowledge regarding PA guidelines and a lack of engagement with content on PA promotion for people with disabilities. The lack of understanding of the PA guidelines and their application to individuals with NC was a barrier, consistent with other studies examining PT knowledge of PA guidelines (Lowe et al., 2017; Rethorn et al., 2021). While PTs are trained in exercise prescription, their training focuses on rehabilitation rather than recreational activity for general health. While there is a call for PTs to support their clients with NC by promoting PA (Quinn & Morgan, 2017), a lack of general knowledge of sport and exercise may hinder PA engagement among clients with NC (Williams et al., 2018). If PTs were better educated on the PA guidelines, PA promotion may become more regular and consistent. Prior research has shown that PT curriculum is being updated to include health promotion strategies (Bodner et al., 2013; Magnusson et al., 2020). How this content is applied, and the effect it has on PA promotion practices, remains to be seen. There have been calls to increase disability-related content in PT training as well; however, large-scale curricular changes are difficult to implement (Garrand et al., 2018).

In the current study, participants reported client-related barriers such as low motivation, client safety, and client finances, and noted that some client factors (such as low motivation) make PTs feel less confident and effective at promoting PA. Some PTs who work with clients with NC have been shown to perceive that their clients' lack of motivation to engage in PA is fixed and that they cannot promote behavior change (Morris et al., 2015; Williams et al., 2018), which is in line with the broader healthcare perspective of promoting behavior change in people who are perceived to have low motivation to engage in PA (Huijg et al., 2015). PTs, like other health professionals, need to be able to address their clients' motivation specifically, even when it is perceived to be low (Yap & Davis, 2008). Perceived low motivation, whether due to a bias on the part of the health professional or as stated by the client, can be addressed.

There have been calls for PTs to enhance their skills in behavior change through motivational interviewing to address patient motivation and adherence (McDevitt

et al., 2024; Pignataro & Huddleston, 2015). It has been shown that motivational interviewing can promote behavior change. A 2011 systematic review and meta-analysis found that motivational interviewing was associated with a greater reduction in body mass compared to controls when promoting weight loss with overweight and/or obese clients (Armstrong et al., 2011). Further, a 2016 study suggested that motivational interviewing can improve health-promoting behaviors in clients with multiple sclerosis (Dashti et al., 2016). Additionally, other behavior change frameworks, such as the Health Belief Model (Rosenstock et al., 1988) and Social Cognitive Theory (Bandura, 1986), can support PTs in addressing perceived low motivation by focusing on self-efficacy, outcome expectations, and barriers to PA engagement (Sweet et al., 2012). PTs can benefit from enhanced education and training to recognize and promote self-esteem, self-efficacy, and positive self-talk in their clients. Addressing perceived low motivation, whether due to a health professional bias or, as the client stated, requires targeted education in evidence-based behavior change techniques such as goal-setting, action planning, and social support development (Michaelsen & Esch, 2023).

This study was not without limitations. PTs included in the study varied in the extent to which they worked with NC. Due to a low response rate, we broadened the criteria to include PTs who work with NC in any capacity. This precludes determining whether the facilitators of and barriers to PA promotion depended on PTs' caseloads. Due to the recruitment method, the current study lacks external validity. Despite these limitations, this study yields important data on this topic.

## Conclusions

This study used the TDF to investigate the barriers to and facilitators of PA promotion by PTs for their clients with NC, to understand why PTs are not consistently promoting PA and to consider implementation strategies to shift this trend. Several barriers and facilitators emerged from this study that are specific to PTs promoting PA for clients with NC, prompting a discussion of strategies to promote facilitators and reduce barriers. The PT profession may benefit from including health promotion and community engagement curricular content, both in student training and in post-professional education. Future research should examine specific implementation strategies to address identified barriers, including enhanced training in behavior change techniques, development of community-clinical partnerships, and organizational supports for PA promotion practices.

## License

This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License.

## References

- Andersen, L. B., Pallesen, T., & Holm Pedersen, L. (2011). Does ownership matter? Public service motivation among physiotherapists in the private and public sectors in Denmark. *Review of Public Personnel Administration, 31*(1), 10–27. <https://doi.org/10.1177/0734371X10394402>
- Anderson, C., Grant, R. L., & Hurley, M. V. (2017). Exercise facilities for neurologically disabled populations – perceptions from the fitness industry. *Disability and Health Journal, 10*(1), 157–162. <https://doi.org/10.1016/j.dhjo.2016.09.006>
- Archibald, M. M., Ambagtsheer, R. C., Casey, M. G., & Lawless, M. (2019). Using Zoom videoconferencing for qualitative data collection: Perceptions and experiences of researchers and participants. *International Journal of Qualitative Methods, 18*, 1609406919874596. <https://doi.org/10.1177/1609406919874596>
- Armstrong, M. J., Mottershead, T. A., Ronksley, P. E., Sigal, R. J., Campbell, T. S., & Hemmelgarn, B. R. (2011). Motivational interviewing to improve weight loss in overweight and/or obese patients: A systematic review and meta-analysis of randomized controlled trials. *Obesity Reviews, 12*(9), 709–723. <https://doi.org/10.1111/j.1467-789X.2011.00892.x>
- Atkins, L., Francis, J., Islam, R., O'Connor, D., Patey, A., Ivers, N., Foy, R., Duncan, E. M., Colquhoun, H., Grimshaw, J. M., Lawton, R., & Michie, S. (2017). A guide to using the Theoretical Domains Framework of behaviour change to investigate implementation problems. *Implementation Science, 12*(1), 77. <https://doi.org/10.1186/s13012-017-0605-9>
- Bandura, A. (1986). *Social foundations of thought and action: A social cognitive theory*. Prentice-Hall, Inc.
- Bodner, M. E., Rhodes, R. E., Miller, W. C., & Dean, E. (2013). Benchmarking curriculum content in entry-level health professional education with special reference to health promotion practice in physical therapy: A multi-institutional international study. *Advances in Health Sciences Education, 18*(4), 645–657. <https://doi.org/10.1007/s10459-012-9404-x>
- Cane, J., O'Connor, D., & Michie, S. (2012). Validation of the theoretical domains framework for use in behaviour change and implementation research. *Implementation Science, 7*(1), 37. <https://doi.org/10.1186/1748-5908-7-37>
- Carroll, D. D., Courtney-Long, E. A., Stevens, A. C., Sloan, M. L., Lullo, C., Visser, S. N., Fox, M. H., Armour, B. S., Campbell, V. A., Brown, D. R., & Dorn, J. M. (2014). Vital signs: Disability and physical activity — United States, 2009–2012. *MMWR. Morbidity and Mortality Weekly Report, 63*(18), 407–413.
- Dashti, A., Yousefi, H., Maghsoudi, J., & Etemadifar, M. (2016). The effects of motivational interviewing on health promoting behaviors of patients with multiple sclerosis. *Iranian Journal of Nursing and Midwifery Research, 21*(6), 640–645. <https://doi.org/10.4103/1735-9066.197682>
- Dean, E. (2009). Physical therapy in the 21st century (Part II): Evidence-based practice within the context of evidence-informed practice. *Physiotherapy Theory and Practice, 25*(5-6), 354–368. <https://doi.org/10.1080/09593980902813416>
- Dzewaltowski, D. A., Estabrooks, P. A., & Glasgow, R. E. (2004). The future of physical activity behavior change research: What is needed to improve translation of research into health promotion practice? *Exercise and Sport Sciences Reviews, 32*(2), 57–63.
- Eccles, M., Grimshaw, J., Walker, A., Johnston, M., & Pitts, N. (2005). Changing the behavior of healthcare professionals: The use of theory in promoting the uptake of research findings. *Journal of Clinical Epidemiology, 58*(2), 107–112. <https://doi.org/10.1016/j.jclinepi.2004.09.002>
- Fereday, J., & Muir-Cochrane, E. (2006). Demonstrating rigor using thematic analysis: A hybrid approach of inductive and deductive coding and theme development. *International Journal of Qualitative Methods, 5*(1), 80–92. <https://doi.org/10.1177/160940690600500107>
- Fernandes, A., Santinha, G., & Forte, T. (2022). Public service motivation and determining factors to attract and retain health professionals in the public sector: A systematic review. *Behavioral Sciences, 12*(4), 95. <https://doi.org/10.3390/bs12040095>
- Francis, J. J., Johnston, M., Robertson, C., Glidewell, L., Entwistle, V., Eccles, M. P., & Grimshaw, J. M. (2010). What is an adequate sample size? Operationalising data saturation for theory-based interview studies. *Psychology & Health, 25*(10), 1229–1245. <https://doi.org/10.1080/08870440903194015>
- French, S. D., Green, S. E., O'Connor, D. A., McKenzie, J. E., Francis, J. J., Michie, S., Buchbinder, R., Schattner, P., Spike, N., & Grimshaw, J. M. (2012). Developing theory-informed behaviour change in-

- terventions to implement evidence into practice: A systematic approach using the Theoretical Domains Framework. *Implementation Science*, 7(1), 38. <https://doi.org/10.1186/1748-5908-7-38>
- Garrand, K., Pechak, C., Jimenez, L., & Gell, N. (2018). Survey of disability competence teaching and assessment strategies in physical therapist education. *Journal of Physical Therapy Education*, 32(1), 55–64. <https://doi.org/10.1097/JTE.0000000000000026>
- Ginis, K. A. M., Ma, J. K., Latimer-Cheung, A. E., & Rimmer, J. H. (2016). A systematic review of review articles addressing factors related to physical activity participation among children and adults with physical disabilities. *Health Psychology Review*, 10(4), 478–494. <https://doi.org/10.1080/17437199.2016.1198240>
- Grimshaw, J. M., Eccles, M. P., Lavis, J. N., Hill, S. J., & Squires, J. E. (2012). Knowledge translation of research findings. *Implementation Science*, 7(1), 50. <https://doi.org/10.1186/1748-5908-7-50>
- Grol, R., & Grimshaw, J. (2003). From best evidence to best practice: Effective implementation of change in patients' care. *Lancet*, 362(9391), 1225–1230. [https://doi.org/10.1016/S0140-6736\(03\)14546-1](https://doi.org/10.1016/S0140-6736(03)14546-1)
- Havercamp, S. M., & Scott, H. M. (2015). National health surveillance of adults with disabilities, adults with intellectual and developmental disabilities, and adults with no disabilities. *Disability and Health Journal*, 8(2), 165–172. <https://doi.org/10.1016/j.dhjo.2014.11.002>
- Huijg, J. M., Gebhardt, W. A., Verheijden, M. W., van der Zouwe, N., de Vries, J. D., Middelkoop, B. J. C., & Crone, M. R. (2015). Factors influencing primary health care professionals' physical activity promotion behaviors: A systematic review. *International Journal of Behavioral Medicine*, 22(1), 32–50. <https://doi.org/10.1007/s12529-014-9398-2>
- Kallio, H., Pietilä, A.-M., Johnson, M., & Kangasniemi, M. (2016). Systematic methodological review: Developing a framework for a qualitative semi-structured interview guide. *Journal of Advanced Nursing*, 72(12), 2954–2965. <https://doi.org/10.1111/jan.13031>
- Kearns, M., Ponichtera, N., Rucker, T., & Ford, G. (2014). Physical therapists as practitioners of choice: Consumer knowledge of practitioner skills and training. *Journal of Physical Therapy Education*, 28(1), 64–72.
- Kennedy, W., Curtin, C., & Bowling, A. (2023). Access to physical activity promotion for people with neurological conditions: Are physical therapists leading the way? *Disability and Health Journal*, 101517. <https://doi.org/10.1016/j.dhjo.2023.101517>
- Kinne, S., Patrick, D. L., & Doyle, D. L. (2004). Prevalence of secondary conditions among people with disabilities. *American Journal of Public Health*, 94(3), 443–445. <https://doi.org/10.2105/AJPH.94.3.443>
- Lobelo, F., Rohm Young, D., Sallis, R., Garber, M. D., Billinger, S. A., Duperly, J., Hutber, A., Pate, R. R., Thomas, R. J., Widlansky, M. E., McConnell, M. V., & Joy, E. A. (2018). Routine assessment and promotion of physical activity in healthcare settings: A scientific statement from the American Heart Association. *Circulation*, 137(18). <https://doi.org/10.1161/CIR.0000000000000559>
- Lowe, A., Littlewood, C., McLean, S., & Kilner, K. (2017). Physiotherapy and physical activity: A cross-sectional survey exploring physical activity promotion, knowledge of physical activity guidelines and the physical activity habits of UK physiotherapists. *BMJ Open Sport & Exercise Medicine*, 3(1), e000290. <https://doi.org/10.1136/bmjsem-2017-000290>
- Magnusson, D. M., Rethorn, Z. D., Bradford, E. H., Maxwell, J., Ingman, M. S., Davenport, T. E., & Bezner, J. R. (2020). Population health, prevention, health promotion, and wellness competencies in physical therapist professional education: Results of a modified Delphi study. *Physical Therapy*, 100(9), 1645–1655.
- McDevitt, A. W., McMullen, J., & Shepherd, M. (2024). Empowering tomorrow's healers: A perspective on integrating person-centered care into physical therapist education. *The Journal of Manual & Manipulative Therapy*, 32(5), 457–463. <https://doi.org/10.1080/10669817.2024.2402100>
- Michaelsen, M. M., & Esch, T. (2023). Understanding health behavior change by motivation and reward mechanisms: A review of the literature. *Frontiers in Behavioral Neuroscience*, 17, 1151918. <https://doi.org/10.3389/fnbeh.2023.1151918>
- Morris, J. H., Oliver, T., Kroll, T., Joice, S., & Williams, B. (2015). From physical and functional to continuity with pre-stroke self and participation in valued activities: A qualitative exploration of stroke survivors', carers' and physiotherapists' perceptions of physical activity after stroke. *Disability and Rehabilitation*, 37(1), 64–77. <https://doi.org/10.3109/09638288.2014.907828>
- Perry, J. L., & Wise, L. R. (1990). The motivational bases of public service. *Public Administration Review*, 50(3), 367–373. <https://doi.org/10.2307/976618>

- Pignataro, R. M., & Huddleston, J. (2015). The use of motivational interviewing in physical therapy education and practice: Empowering patients through effective self-management. *Journal of Physical Therapy Education*, *29*(2), 62–71. <https://doi.org/10.1097/00001416-201529020-00009>
- Quinn, L., & Morgan, D. (2017). From disease to health: Physical therapy health promotion practices for secondary prevention in adult and pediatric neurologic populations. *Journal of Neurologic Physical Therapy*, *41*, S46–S54. <https://doi.org/10.1097/NPT.000000000000166>
- Ramanadhan, S., Revette, A. C., Lee, R. M., & Aveling, E.-L. (2021). Pragmatic approaches to analyzing qualitative data for implementation science: An introduction. *Implementation Science Communications*, *2*(1), 70. <https://doi.org/10.1186/s43058-021-00174-1>
- Rethorn, Z. D., Covington, J. K., Cook, C. E., & Bezner, J. R. (2021). Physical activity promotion attitudes and practices among outpatient physical therapists: Results of a national survey. *Journal of Geriatric Physical Therapy*, *44*(1), 25–34. <https://doi.org/10.1519/JPT.0000000000000289>
- Rimmer, J. H. (2012). Getting beyond the plateau: Bridging the gap between rehabilitation and community-based exercise. *PM&R*, *4*(11), 857–861. <https://doi.org/10.1016/j.pmrj.2012.08.008>
- Rimmer, J. H., Chen, M.-D., & Hsieh, K. (2011). A conceptual model for identifying, preventing, and managing secondary conditions in people with disabilities. *Physical Therapy*, *91*(12), 1728–1739. <https://doi.org/10.2522/ptj.20100410>
- Rimmer, J. H., Vanderbom, K. A., Bandini, L. G., Drum, C. E., Luken, K., Suarez-Balcazar, Y., & Graham, I. D. (2014). GRAIDs: A framework for closing the gap in the availability of health promotion programs and interventions for people with disabilities. *Implementation Science*, *9*(1), 100. <https://doi.org/10.1186/s13012-014-0100-5>
- Robey, K. L., Minihan, P. M., Long-Bellil, L. M., Hahn, J. E., Reiss, J. G., & Eddey, G. E. (2013). Teaching health care students about disability within a cultural competency context. *Disability and Health Journal*, *6*(4), 271–279. <https://doi.org/10.1016/j.dhjo.2013.05.002>
- Roh, C.-Y., Moon, M. J., Yang, S.-B., & Jung, K. (2016). Linking emotional labor, public service motivation, and job satisfaction: Social workers in health care settings. *Social Work in Public Health*, *31*(2), 43–57. <https://doi.org/10.1080/19371918.2015.1087904>
- Rosenstock, I. M., Strecher, V. J., & Becker, M. H. (1988). Social learning theory and the Health Belief Model. *Health Education Quarterly*, *15*(2), 175–183. <https://doi.org/10.1177/109019818801500203>
- Ross, S. M., Bogart, K. R., Smit, E., Hatfield, B., Yun, J., & Logan, S. W. (2021). Physical activity, medical home, and health behavior counseling among adolescents with special health care needs: NSCH 2016–2017. *Maternal and Child Health Journal*, *25*(4), 542–553. <https://doi.org/10.1007/s10995-020-03089-w>
- Sanchez, A., Bully, P., Martinez, C., & Grandes, G. (2015). Effectiveness of physical activity promotion interventions in primary care: A review of reviews. *Preventive Medicine*, *76*, S56–S67. <https://doi.org/10.1016/j.ypmed.2014.09.012>
- Sandelowski, M. (2000). Whatever happened to qualitative description? *Research in Nursing & Health*, *23*(4), 334–340. [https://doi.org/10.1002/1098-240X\(200008\)23:4%3C334::AID-NUR9%3E3.0.CO;2-G](https://doi.org/10.1002/1098-240X(200008)23:4%3C334::AID-NUR9%3E3.0.CO;2-G)
- Sharp, A., & Herrman, D. (2021). Disability and physical therapy: A complicated relationship, an uncertain path forward. *Physical Therapy*, *101*(7), pzab085. <https://doi.org/10.1093/ptj/pzab085>
- Sumathi, G. N., Kamalanabhan, T. J., & Thenmozhi, M. (2015). Impact of work experiences on perceived organizational support: A study among healthcare professionals. *AI & Society*, *30*(2), 261–270. <https://doi.org/10.1007/s00146-013-0509-4>
- Sweet, S. N., Fortier, M. S., Strachan, S. M., & Blanchard, C. M. (2012). Testing and integrating self-determination theory and self-efficacy theory in a physical activity context. *Canadian Psychology / Psychologie Canadienne*, *53*(4), 319–327. <https://doi.org/10.1037/a0030280>
- Taukobong, N. P., Myezwa, H., Pengpid, S., & Van Geertruyden, J.-P. (2014). The degree to which physiotherapy literature includes physical activity as a component of health promotion in practice and entry level education: A scoping systematic review. *Physiotherapy Theory and Practice*, *30*(1), 12–19. <https://doi.org/10.3109/09593985.2013.783896>
- Tranter, K. E., Harvey, L. A., Chu, J., Ilha, J., Ben, M., Chen, L. W., & Glinsky, J. V. (2025). Using the theoretical domains framework to identify strategies to support the implementation of the guidelines for the physiotherapy management of people with spinal cord injury: A qualitative study. *Spinal Cord Series and Cases*, *11*(1), 25. <https://doi.org/10.1038/s41394-025-00719-9>
- van der Ploeg, H. P., Streppel, K. R. M., van der Beek, A. J., van der Woude, L. H. V., Vollenbroek-Hutten, M. M. R., van Harten, W. H., & van

- Mechelen, W. (2007). Successfully improving physical activity behavior after rehabilitation. *American Journal of Health Promotion, 21*(3), 153–159. <https://doi.org/10.4278/0890-1171-21.3.153>
- Verhagen, E., & Engbers, L. (2008). The physical therapist's role in physical activity promotion. *British Journal of Sports Medicine, 43*(2), 99–101. <https://doi.org/10.1136/bjsm.2008.053801>
- Whalen Smith, C. N., Havercamp, S. M., Tosun, L., Shetterly, S., Munir, A., Kennedy, W., Feldner, H. A., Herrman, D., Sloane, B. M., & Weinstein, F. H. (2024). Training an anti-ableist physical therapist workforce: Critical perspectives of health care education that contribute to health inequities for people with disabilities. *Physical Therapy, 104*(9), pzae092. <https://doi.org/10.1093/ptj/pzae092>
- Williams, T. L., Smith, B., & Papathomas, A. (2018). Physical activity promotion for people with spinal cord injury: Physiotherapists' beliefs and actions. *Disability and Rehabilitation, 40*(1), 52–61. <https://doi.org/10.1080/09638288.2016.1242176>
- Yap, T. L., & Davis, L. S. (2008). Physical activity: The science of health promotion through tailored messages. *Rehabilitation Nursing, 33*(2), 55–62. <https://doi.org/10.1002/j.2048-7940.2008.tb00204.x>

## Appendix

**Table 2: Facilitators**

**Table 3: Barriers**

**Table 4: Semi-Structured Interview Guide**

**Table 2:** *Facilitators*

Facilitator	Factor	Exemplar Quotes
Personal	High motivation and confidence in promoting PA for patients with NC	“I have no problem [confidence in promoting PA]. I kind of figured the neuromuscular disorder is just another challenge that they have to meet. And if nothing else they’re going to benefit so much more from maintaining or gaining strength and more fitness than just a general population.” [P4]
Personal	A sense of responsibility to learn about PA promotion	“[Response to question asking what would facilitate learning about PA promotion for people with ND] It would be just kind of on my own trying my best to kind of keep up.” [P1]
Personal	A desire to model PA behaviors in order to better understand and support the experiences of their patients	“If I’m providing a guideline or, recommending that somebody become more physically active and I don’t do it myself, I feel, I’m not practicing what I’m preaching, so I feel like it gives me a certain level of, not respect, but I know what I’m talking about because I do it myself.” [P6]
Personal	Personal enjoyment/satisfaction promoting PA to patients	“It’s very satisfying when you can motivate somebody to do it, to participate in physical activity that they maybe weren’t doing before. I think that can be very rewarding personally.” [P10]
Professional	Connections to community PA opportunities that they promote to their patients with NC	“I have an in at the university. My husband is the disability accommodation director at the state university here, so I pick his brain because he has resources at the university. . . . Also, there’s Special Olympics teams here, so I’m picking their brains to see.” [P8]
Professional	Availability of continuing education relevant to PA promotion for patients with ND	“I took some courses related to physical activity and prevention of obesity for people with disabilities and the prevalence is so much greater because they don’t sometimes get that physical activity. I took a number of courses related specifically to that, and I really loved them.” [P3]
Professional	PT education had relevant content related to people with disabilities	“I think that it was probably separated out into a class related to disability studies and a little bit more of the awareness of assistive devices and things that they might need for their daily activity. And then the recommendations for physical activity across the life and age spectrum with considerations.” [P7]
Professional	Job setting/environment facilitates PA promotion	“The job location that I have right now participates in silver sneakers. so, we do have that ability for people to come in and use our facility to work out, so that makes it a lot easier for me to have somebody who comes in that I think could benefit from health promotion. . . . I like the fact that our facility is open and available to encourage patients to come back to it.” [P4]
Client-Related	Highly motivated	“Some people are very enthusiastic about kind of taking care of their lives, changing their lifestyles, kind of into healthy living and healthy kind of activities. So, these are something that when they’re motivated to do it. . . that is encouraging and then you give them more opportunities to get them the options to do it.” [P1]

**Table 3:** *Barriers*

Barrier	Factor	Exemplar Quotes
Professional	Lack of specific knowledge on PA guidelines	“I don’t know of it [PA guidelines] and it’s probably crap.” [P5]
Professional	Lack of content on PA promotion for people with disabilities in PT training	“I don’t think that that was considered [PA promotion for people with disabilities]. I have been a long-time graduate, so it’s been a while since I graduated, and I don’t think that at that time physical activity was kind of considered a part of it.” [P1]
Professional	Lack of specific continuing education resources on PA promotion	“I’ve attended courses at CSM [combined sections meeting through APTA] that are related to health promotion and then done a lot of reading on my own. I don’t know that I’ve attended any specific courses about health promotion [for people with disabilities] though.” [P10]
Professional	Time	“If I had more ability to have more time with my patients, that would help me, but, you know, limited time. . . . time would be nice.” [P6]
Professional	Cost as a barrier to learn about PA promotion	“I mean most PTs who halfway like their job would take a continuing education course every weekend, but I don’t always have \$600 to just be like here.” [P5]
Professional	Lack of consistency and organizational support and/or guidelines for PA promotion within the profession	“I think we’re not given like a lot of guidelines on what that [PA promotion] should be. . . . that’s up to our just own discretion as a therapist.” [P2]
Professional	Billing/insurance	“Yeah, so for example, payment barriers. That would be big part of it. . . . I think that part of the barrier is like a push for productivity as far as like a push for trying to make sure we bill the maximum number of units that we can.” [P4]
Client-Related	Low patient motivation	“There are people who don’t want to do it [PA]. They don’t like the way they feel when they move. And that becomes very challenging. Finding a place of entry is hard. . . . it’s really hard to convince one person and a family to do it when no one else is.” [P7]
Client-Related	Patient safety	“They are going to be the same barriers [when comparing patients with and without NC], but with a few added ones and that is, do I have people around me who can help when I don’t need help but still be there? Am I safe? Where am I going to do my workouts? Even at home, am I going to be okay if I fall down? . . . So, I think there’s another layer [which is] safety.” [P8]
Client-Related	Financial and insurance resources	“I don’t think I would bring up a ton of activities to parents. If I knew they cost money for them. I try to only tell them things that are free.” [P2]

**Table 4:** *Semi-Structured Interview Guide Informed by the TDF*

Construct	Definition	Question	Prompts
Knowledge	An awareness of the existence of something.	What is your current understanding of health-promoting physical activity guidelines?	What is your current understanding of how they should be modified for individuals with neuromuscular disorders (NMD)?
Skills	An ability or proficiency acquired through practice.	What were the courses you took in PT school that focused on health enhancing PA promotion for people with disabilities?	Have you pursued any continuing education in this area? Have you had opportunity to?
Social/Professional Role and Identity	A coherent set of behaviors and displayed personal qualities of an individual in a social or work setting.	What role do you think PTs should play in health enhancing PA promotion for individuals with NMD?	
Beliefs about Capabilities	Acceptance of the truth, reality, or validity about an ability, talent, or facility that a person can put to constructive use.	How confident do you feel in promoting health enhancing PA for individuals with NMD?	For each PA promotion type: how confident do you feel? Give examples of you doing the ones that you are confident in.
Optimism	The confidence that things will happen for the best or that desired goals will be attained.	How effective do you think your health enhancing PA promotion efforts are for this population?	Are there things you can do better? What are some barriers that your patients face?
Beliefs about Consequences	Acceptance of the truth about outcomes of a behavior in a given situation.	What are some of the benefits of health-enhancing PA for folks with mobility disabilities?	
Intentions	A conscious decision to perform a behavior or a resolve to act in a certain way.	What influences your decision to promote health enhancing PA to one of your patients with NMD?	
Goals	Mental representations of outcomes or end states that an individual wants to achieve.	Do you believe the health enhancing PA guidelines are a good goal for individuals with NMD?	
Innovation	Any characteristics of the innovation that discourages or encourages the development of skills and abilities.	What are some of the facilitators that might make it easier to promote health enhancing PA?	What are some barriers facing PTs promoting health enhancing PA for their patients with NMD?
Socio-political Context	Any characteristics of the socio-political context that discourages or encourages development.	What are financial facilitators or barriers to including health enhancing PA promotion in treatment sessions?	Can you bill for this? Do you bill for this?
Organization	Any characteristics of the organization that discourages or encourages development.	In what ways does your job setting encourage you to promote health enhancing PA for individuals with NMD?	Are there any employer facilitators or barriers?
Patient	Any characteristics of the patient that discourages or encourages development.	What are some patient characteristics that may deter you from promoting PA?	What are some patient characteristics that may facilitate you promoting PA?
Innovation Strategy	Any characteristics of the innovation strategy that discourages or encourages development.	What are facilitators to you learning more about health enhancing PA promotion for people with NMD?	What are barriers to learning more?

*Table 4 continued*

<b>Construct</b>	<b>Definition</b>	<b>Question</b>	<b>Prompts</b>
Social Influences	Those interpersonal processes that can cause individuals to change their thoughts, feelings, or behaviors.	What are some things your colleagues do to promote PA for individuals with NMD?	What do you think APTA does or could do better to help you promote health enhancing PA?
Positive Emotion (Motivation)	A complex positive reaction pattern by which the individual attempts to deal with a personally significant matter.	What benefits do you get from promoting health enhancing PA for individuals with NMD?	
Negative Emotions	A complex negative reaction pattern by which the individual attempts to deal with a personally significant matter.	What are some drawbacks or frustrations to promoting health enhancing PA for individuals with NMD?	
Behavioral Regulation	Anything aimed at managing or changing objectively observed or measured actions.	Is PA a regular part of your life?	Do you find it important to meet physical activity guidelines? Explain.
Nature of the Behaviors	The nature of the aggregate of all responses made by an individual in any situation.	How hard do you think it is to include health enhancing PA promotion in conjunction with your treatment plan?	
Knowledge/Environment	Awareness of resources available to implement behavior.	Are you aware of any resources or facilities that help individuals with NMD engage in PA in your community?	
Motivation	The individual's level of wanting to engage in the behavior.	How motivated are you to promote health enhancing PA for individuals with NMD?	Do you experience competing priorities for your time that you are more motivated to meet? Explain.